



Information for Immunizers

COMIRNATY® COVID-19 mRNA vaccine

12 years of age and older


- COMIRNATY® is administered intramuscularly as a single dose of 0.3 mL, regardless of prior COVID-19 vaccination status.
- For individuals who have previously been vaccinated with a COVID-19 vaccine, COMIRNATY should be administered at least 3-6 months after the most recent dose of a COVID-19 vaccine.

Vaccine is provided in a Multiple Dose Vial: **DO NOT DILUTE**

COMIRNATY® multiple dose vial with a *gray cap and a gray label* border does not contain preservative.

COMIRNATY® is intended for individuals 12 years of age and older (gray cap/gray label border) and cannot be used to prepare doses for individuals aged 6 months to <12 years of age.

- Please ensure each vial is thawed prior to administration. Reminder **DO NOT DILUTE** prior to use.
- Thawed vials can be stored in the refrigerator (2°C to 8°C) for up to 10 weeks prior to use within the expiry date printed on the carton.
- Thawed vials can be stored at room temperature (up to 25°C) for up to 12 hours prior to use.
- The thawed suspension may contain white to off-white opaque amorphous particles.
- After first puncture, record the time, store between 2°C to 25°C and use within 12 hours.
- One multiple dose vial contains up to 6 doses of 0.3 mL.
- Low dead-volume syringes and/or needles can be used to extract 6 doses from a single vial. In order to ensure consistent withdrawal of 6 doses of 0.3 mL, it is important to adhere to minimizing volume loss during dose extraction.
- If the amount of vaccine remaining in the vial cannot provide a full dose of 0.3 mL, discard the vial and any excess volume.



Vials must be discarded 12 hours after first puncture

Full Product Monograph available online: [comirnaty-kp2-pm-en.pdf \(canada.ca\)](https://www.canada.ca/comirnaty-kp2-pm-en.pdf)